

Self-reported Dominance in Women: Associations with Hormonal Contraceptive use, Relationship Status, and Testosterone

Kelly D. Cobey · Mike Nicholls ·
Juan David Leongómez · S. Craig Roberts

Received: 21 October 2014 / Revised: 10 February 2015 / Accepted: 13 February 2015 /
Published online: 10 March 2015
© Springer International Publishing 2015

Abstract How to achieve dominance in a group is a recurrent challenge for individuals of many species, including humans. Previous research indicates that both relationship status and contraceptive use appear to moderate women's testosterone levels. If testosterone contributes to dominance, this raises the possibility for group differences in dominance between single and partnered women, and between users and non-users of hormonal contraception. Here, we examine associations between relationship status and use/non-use of hormonal contraception and women's self-reported social dominance. In a sample of 84 women, we replicate previous research documenting a significant positive correlation between women's saliva testosterone levels and their self-reported dominance. Consistent with other literature, we also find that women using hormonal contraception have significantly lower testosterone than those who are regularly cycling and that partnered women have significantly lower testosterone than single women. Although we do not find a main effect of either relationship status or hormonal contraceptive use status on women's reported levels of dominance, the interaction between these variables predicted reported dominance scores. This interaction remained significant when participant age and testosterone values were added to the model as covariates. We discuss these results in the context of the existing literature on testosterone and women's dominance behaviour and with respect to the evolutionary benefits of social dominance in women.

K. D. Cobey (✉) · M. Nicholls · J. D. Leongómez · S. C. Roberts
Division of Psychology, University of Stirling, Stirling, United Kingdom
e-mail: Kelly.Cobey@stir.ac.uk

K. D. Cobey
Laboratory of Experimental and Comparative Ethology (LEEC), University of Paris 13, Paris, France

J. D. Leongómez
Facultad de Psicología, Universidad El Bosque, Bogotá, Colombia

Keywords Testosterone · Dominance · Hormonal contraception · Relationship status · Self-reported Dominance

Introduction

Dominance within a hierarchy is often associated with greater access to resources and with a greater ability to influence subordinates (Magee and Galinsky 2008). A distinction can be made between social dominance, wherein an individual is interpersonally dominant, and dominance rank, one's social standing within a group. Given the potential adaptive benefits of dominance, considerable effort has been made to understand the underlying mechanisms that regulate the expression of both of these dominance forms. In particular, previous research has considered the potential for a relationship between dominance and testosterone levels. The literature on this topic generally shows a consistent picture when considering non-human animals. For example, correlational studies in male primates examining dominance rank have shown that high-ranking individuals produce higher levels of testosterone than their lower ranked counterparts (e.g. Eastern common chimpanzee, *Pan troglodytes schweinfurthii*: Muller and Wrangham 2004; Rhesus monkey, *Macaca mulatta*: Rose, Holaday, and Bernstein 1971). Similar correlational effects between testosterone and dominance have been documented among a range of other species including birds (Harding 1983), and hamsters, dogs and deer (Rada, Kellner, and Winslow 1976). Work which has experimentally manipulated testosterone, and subsequently measured shifts in dominance, appears to confirm these associations. For example, testosterone reduction results in a loss of social dominance in rats (Albert, Walsh, Gorzalka, Siemens, and Louie 1986), and implanting or injecting testosterone increases the expression of social dominance in a number of other species (e.g. Bouissou 1990; Searcy and Wingfield 1980). Among human males, there is also relatively consistent evidence for a positive relationship between testosterone and social dominance (e.g. Mehta and Josephs 2010; Neave, Laing, Fink, and Manning 2003; Swaddle and Reiersson 2002).

By comparison, the study of dominance in females has received far less attention than it has in males. This is regrettable since females, like males, can benefit from being dominant in a number of different contexts. For example, females can use social dominance to compete more readily with other females to attract mates and in order to gather resources necessary to achieve status or to stay healthy (Darwin 1871). Among human females, in contrast to the work in males, the evidence for a relationship between testosterone and dominance is comparatively mixed and, in many cases, it is only after the consideration of a moderating variable that correlations are found.

Early work examining the relationship between testosterone and dominance in women produced positive correlations. For example, Purifoy and Koopmans (1979) found that 'career-oriented' women (those who had professional careers, technical jobs, or worked in management) tended to have higher testosterone levels than women who did not work or who worked in lower-level clerical roles. In this study, career-orientation was argued to reflect traits such as 'assertiveness', which is considered to be a component of socially dominant behaviour, thereby suggesting a link between testosterone and dominant career traits. Likewise, in a study where women self-reported the extent to which a series of adjectives applied to them, those with higher testosterone levels identified as being more

‘dominant’ (Udry and Talbert 1988). Furthermore, Grant and France (2001) reported that women’s self-reported dominance was positively associated with testosterone levels.

Work that built on these initial studies produced more varied results. For example, Edwards and Casto (2013) found a relationship between dominance and testosterone, but only in women with relatively low cortisol. Mehta and Josephs (2010) documented a similar moderating effect of cortisol on the relationship between testosterone and social dominance in their earlier work. In contrast, Denson, Mehta, and Ho Tan (2013) found the opposite effect, namely that testosterone and dominance were positively related, but only among women with high cortisol. As a result, while there is evidence that cortisol may moderate the association between testosterone and dominance, the direction of this effect is inconsistent across studies. Furthermore, other studies have documented null effects for the relationship between testosterone and dominance in women (Kivlighan et al. 2005; Stanton and Schultheiss 2007) and one study (Cashdan 1995) actually found a negative correlation between social rank and androgen levels in women. These somewhat mixed results suggest that the association between dominance and testosterone in women may be more variable than it is among men, and may perhaps be moderated by other socially relevant factors.

Here, we seek to examine how two social factors, namely relationship status and hormonal contraceptive use, might influence women’s social dominance. Both relationship status and use of hormonal contraception are known to influence testosterone levels, meaning that there could be group differences in social dominance among single and partnered women, and among users and non-users of hormonal contraception. Research examining the impact of relationship status on endocrine variation suggests that paired individuals have lower testosterone levels than single individuals (e.g. in men, Burnham et al. 2003; in women, Kuzawa, Gettler, Huang, and McDade 2010; in heterosexual men and homosexual women, van Anders and Watson 2006). There is some additional evidence that this effect of relationship is moderated differently in men and women. For example, van Anders and Goldey (2010) found that the effect in women was moderated by frequency of sexual activity, while the effect in men was moderated by interest in new partners. Interestingly, however, there is no evidence that being the dominant versus subordinate individual in a couple relates to measured cortisol (Maestripieri et al. 2013). This suggests that relationship status is a better predictor of hormone levels than relationship dominance status. With respect to the influence of hormonal contraceptive use on women’s hormone levels, prospective studies demonstrate that women exhibit a marked reduction in testosterone following initiation of hormonal contraceptive use (Zimmerman, Eijkemans, Coelingh Bennink, Blankensein, and Fauser 2013a; Zimmerman et al. 2014).

Dominance is no doubt the product of a series of complex physiological and cognitive components, meaning that there is likely a range of variables that potentially influence its expression. In spite of this, to our knowledge, no previous study examining women’s social dominance has considered, or controlled for, both relationship status and contraceptive pill use in the same experiment. This omission is somewhat surprising given the literature suggesting that both variables influence women’s testosterone levels. For these reasons, this study was designed to test the following predictions: (1) Self-reported dominance in women will be positively related to testosterone levels; (2) Women using hormonal contraception will report lower levels of dominance than regularly cycling women; (3) Partnered women will report lower levels of dominance than single women.

Methods

Participants

Participants were 85 women who were recruited from the University of Stirling. All procedures were approved by the Psychology Departmental Ethics Review Board. Participants were aged between 18 and 32 years (Mean=21.05, S.D. = 3.17, $N=7$ missing). Of this sample, 29 women reported to be using some form of hormonal contraception, while 56 reported to have regular menstrual cycles (cycles between 25–35 days in length). Forty-seven of the participants reported to be in a relationship (32 contraceptive users, 15 regularly cycling women), while 38 indicated they were single (11 contraceptive users, 27 regularly cycling women). Of the overall group, eight identified as being non-heterosexual. The final analysis included 82 women. One participant was excluded because her saliva sample was insufficient for assay, one because her testosterone values were greater than 2SD above the mean, and one because she failed to complete the questionnaire measures in their entirety.

Procedure

In the recruitment material for the study it was made clear to participants that they should not eat or drink within 1 h prior to the test session. Moreover, participants were asked to avoid consuming alcohol for 24 hours before participation and not to take part in vigorous exercise on the day of testing, because these factors can impact hormone levels (e.g. Gill 2000; Kraemer and Ratamess 2005). Experimental sessions took place between 1:00pm and 6:00pm. This window was chosen to help control for diurnal rhythms in testosterone levels (e.g. Dabbs 1990). Upon reporting to the lab, participants provided informed consent and were then instructed to rinse their mouth with water. Before any questionnaire items were completed, and after a rest period, participants provided a saliva sample that was collected in 2ml cryovials via passive drool. Thereafter, participants completed an online questionnaire. This questionnaire contained basic demographic items and the IPIP scale for social dominance, which contains 11 items and is designed to measure individual differences in dominance (<http://ipip.ori.org/newIndexofScaleLabels.htm>). This measure has been used in a wide range of previous studies and in our study it had good reliability (Cronbach's Alpha=.83). To complete this scale, participants read a series of statements and then indicated, on a scale of one to seven, the extent to which they agree with each. Higher scores indicate higher levels of agreement. Examples of scale items include 'I am not afraid of providing criticism' and 'I am quick to correct others'. Participants also self-identified as being 'partnered' or not, we did not impose a minimum duration of relationship to be considered partnered.

Hormonal assays

Saliva samples were immediately frozen and stored at -20°C following collection. The samples were then collectively shipped on dry ice to Salimetrics UK Laboratory (Suffolk, UK) where they were analysed for testosterone in duplicate using enzyme immunoassay. On the day of assay, samples were completely thawed and then vortexed and centrifuged at 3000rpm for 15 min. Salimetrics used 25 μl of saliva per

determination with a lower limit of sensitivity of less than 1.0 pg/ml. The intra-assay coefficients of variation were between 2.5% and 6.7%. Previous research indicates that testosterone measures obtained from saliva are positively and strongly related to those obtained via serum samples (Johnson, Joplin, and Burrin 1987; Vittek, L’Hommedieu, Gordon, Rappaport, and Louis Southren 1985).

Results

Descriptive analyses

The mean testosterone level among our 82 included participants was 72.21 pg/mL, with values ranging from 27.99 pg/ml to 141.27 pm/mL (SD=26.29 pg/ml). These values are similar to those reported in previous research using a similar sample (e.g. Deady, Smith, Sharp, and Al-Dujaili 2006; Welling et al. 2007). Spearman’s rank correlations indicated that neither the time of sampling ($r=-.01$, $p=.92$) nor participants’ age ($r=-.14$, $p=.23$) were related to measured testosterone levels. This is likely due to the relatively restrictive timeframe in which we collected our samples and the high degree of homogeneity in this sample of women.

The relationship between testosterone and social dominance

We conducted a one-tailed Spearman’s correlation to examine the relationship between testosterone and women’s self-reported social dominance. The results of this directional test indicated a weak positive correlation ($r_s=.20$, $p=.04$; Fig. 1). A partial correlation controlling for participant age, participant relationship status and participant use/non-use of hormonal contraception remained significant ($r_s=.023$, $p=.03$).

Effects of relationship status and contraceptive pill use on social dominance

To determine how relationship status and hormonal contraceptive use influenced self-reports of dominance, we conducted an analysis of variance with relationship status and




Fig. 1 The relationship between testosterone levels and self-reported social dominance levels in women

hormonal contraceptive use as fixed-factors and dominance scores on the IPIP scale as the dependent variable. This allowed us to look for group differences between users and non-users of hormonal contraception and single and partnered women. We added testosterone levels and participant age as covariates.

We found no main effect of either relationship status ($F=1.17, p=.28$) or hormonal contraceptive pill use ($F=.40, p=.53$) in this initial model. However, the interaction of these two variables on social dominance was statistically significant ($F=6.49, p=.01$; Fig. 2), such that, among regularly cycling women, partnered individuals had significantly higher social dominance scores ($M=43.63, S.E. = 2.49$) than those who were single ($M=34.89, S.E. = 2.03$) (post hoc $t=2.09, p=.04$), while among hormonal contraceptive users, single women had higher levels of social dominance ($M=42.60, S.E. = 3.15$) than partnered women ($M=39.03, S.E. = 1.79$) (post hoc $t=.90, p>0.05$). Testosterone levels also proved to be a significant predictor in this model ($F=5.43, p=.02$), but participant age did not have a significant effect ($F<.01, p=.97$). The results of this model were unchanged when testosterone and participant age were removed as well as when we restricted our analyses to women using combined oral contraceptives (e.g. excluding users of other forms of HC) and when we excluded non-heterosexual women ($N=8$).

Note also that a univariate ANOVA comparing levels of testosterone among users and non-users of hormonal contraception was significant ($F=9.21, p=.003$), with users having lower levels than non-users as is consistent with previous research (e.g. Zimmerman et al. 2013b) (Fig. 3). In spite of this difference, users and non-users did not differ in their levels of social dominance ($F=.07, p=.79$). Likewise, women who were partnered had significantly lower testosterone levels ($M=66.38, S.E. = 3.77$) than those who were single ($M=79.67, S.E. = 2.47$) ($F=5.45, p=.02$); but the groups did not differ in social dominance ($F=1.24, p=.27$).

Discussion

Our results indicate that, in this sample, women's measured testosterone levels show a positive but weak relationship with their self-reported dominance scores. Although the




Fig. 2 Social dominance for hormonal contraceptive (HC) users and non-users, split by relationship status (single: white bars; partnered: grey bars). Bars represent mean \pm 1 s.e.m. For post-hoc tests (pairwise comparisons) the dashed line represents the comparison between partnered HC users and non-users, and the dotted line represents the comparison between single HC users and non-users; † $p<0.1$, * $p<0.05$, ** $p<0.01$




Fig. 3 Levels of testosterone among hormonal contraceptive (HC) users were significantly lower than among women who were regularly cycling. $**p < 0.01$

main effects of relationship status and hormonal contraceptive use on levels of dominance were not significant, suggesting the absence of overall group differences in social dominance among single and partnered women or among hormonal contraceptive users and non-users, we find that the interaction between these two factors does predict women's self-reported dominance scores. Thus, our results suggest that while testosterone and dominance in women may be positively related, dominance is also affected by the interaction of a woman's current hormonal contraceptive use and her relationship status. More broadly put, our findings suggest that women's dominance, an important tool to achieve status, varies with current social circumstances.

When examining the documented interaction in more detail, our results indicate that, among regularly cycling women, partnered women tended to have lower testosterone levels than those who were single, but were significantly more dominant. This pattern of results was in contrast to our prediction that partnered women would have lower levels of dominance than single women as a result of lowered testosterone levels. Our prediction was based on a series of studies examining testosterone levels in single and partnered individuals, which show that the former tend to have significantly higher testosterone values, arguably as a result of the greater relative importance of mating effort for this group (Archer 2006; Wingfield et al. 1990). Our finding that single individuals have higher testosterone is consistent with results found in several independent labs and across cultures (Burnham et al. 2003; Gray et al. 2004; McIntyre et al. 2006; van Anders and Watson 2006, 2007). One interesting point with respect to previous research on testosterone and relationship status is that the vast majority have tested only male participants. A few studies do show analogous effects in women (e.g. van Anders and Watson 2006), although this effect was only present in non-heterosexual women. Other work has documented null or mixed effects in men as well as in women (e.g. McIntyre et al. 2006; Pollet, van der Meij, Cobey, and Buunk 2011). Our findings suggest that while testosterone is higher in single women who are regularly cycling compared to those who are partnered, this does not lead to differences in levels of social dominance between the groups. It may therefore be that high testosterone is a necessary, but not sufficient, component for the expression of women's social dominance.

Moreover, the context in which social dominance is expressed may have important implications for the influence of both social and biological variables. The scale we used

to assay dominance was very general, meaning future research could examine how dominance in different domains (e.g. mating versus friendship) is related to variables like hormonal contraceptive use and relationship status, and how this behaviour relates to measured testosterone. Indeed, dominance may be context-dependent or independent and this was not appropriately considered in the current work. Moreover, the extent to which our self-reported measures of dominance reflect personality traits as apposed to a manifestation of power (reviewed in Ponzi et al. 2015) remains unclear. Our measure of dominance is likely to have conflated social dominance and dominance rank; refined work that looks at dominance in social and hierarchical settings is therefore needed. Future research could similarly build on the existing findings by measuring women's dominance in a behavioural task rather than via self-report questionnaires.

One explanation for our finding that partnered women are more socially dominant may be that the nature of the relationships among the women in our sample may differ from those used in previous studies. For example, relationships may be of differing durations and levels of commitment, or women may differ in levels of sociosexuality. Indeed, several of these factors have previously been examined in association to relationship status and testosterone, but the effects appear to be mixed (Edelstein, Chopik, and Kean 2011; McIntyre et al. 2006; van Anders and Goldey 2010; van Anders and Watson 2007). An alternative explanation may be that relatively dominant women are more likely to seek out and successfully obtain a relationship. Prospective studies which track women's social dominance as they enter and leave relationships will be important to verify effects documented herein.

With respect to the impact of hormonal contraceptive use on women's endocrine levels, research indicates that use of hormonal contraception results in a marked reduction in testosterone levels (e.g. Zimmerman et al. 2013). This was also the case in our study, with users of hormonal contraception having significantly lower levels of testosterone than non-users. Based on this difference, we predicted that women using hormonal contraception might display lower levels of dominance, but we found no main effect of hormonal contraceptive use in our study. This might suggest that a series of complex physiological and cognitive components, beyond solely testosterone levels, combine to influence social dominance levels. Interestingly, in previous work we have shown prospectively that partnered women who initiate hormonal contraception experience a decrease in their levels of intrasexual (female-female) competition (Cobey, Klipping, and Buunk 2013). This may again suggest that the context of social dominance should be specified in future research. Although we attempted to control for the type and method of administration of hormonal contraception by conducting subsequent analyses in our sample, future research using larger samples could consider the impact on testosterone levels and dominance of specific hormonal contraceptive doses and variants of synthetic estrogen and progesterone (see Cobey, Pollet, Roberts, and Buunk 2011; Piccoli, Cobey, and Carnaghi 2014; Welling, Puts, Roberts, Little, and Burriss 2012). Likewise, further studies could consider the influence of cycle phase in regularly cycling women, as there is some evidence to suggest that testosterone increases near to ovulation (e.g. Alexander, Sherwin, Bancroft, and Davidson 1990; Bloch, Schmidt, Su, Tobin, and Rubinow 1998; Dabbs 1990; Morris, Udry, Khan-Dawood, and Dawood 1987; Schultheiss, Dargel, and Rohde 2003; Welling et al. 2007, but see Dabbs 1990; Liening, Stanton, Saini, and Schultheiss 2010; Schultheiss et al. 2003), although this effect is unlikely to be large since daily fluctuations (due to

circadian rhythm) and seasonal fluctuations in testosterone are often greater than cyclical changes.

The critical question of what mechanism (s) underlies dominance remains. One limitation of the current work is that it used a single assay of women's testosterone levels that was taken in the late afternoon. Future research utilizing a larger range of hormone assays and multiple samples and including, for example, measurements of waking testosterone levels, are thus warranted. While our sample size is modest, our results emphasise the idea that while dominance among women may be correlated with testosterone level, and thus is biologically based, its expression is also contingent on social variables. Understanding whether and how hormonal contraception influences, or interacts to influence, dominance levels is potentially important for women's ability to make an informed decision about using this modern intervention (Cobey and Buunk 2012; Roberts, Cobey, Klapilová, and Havlíček 2014). We hope that our results stimulate further studies that consider a broader range of social variables that may impact the expression of women's social dominance.

References

- Albert, D. J., Walsh, M. L., Gorzalka, B. B., Siemens, Y., & Louie, H. (1986). Testosterone removal in rats results in a decrease in social aggression and a loss of social dominance. *Physiology & Behavior*, *36*, 401–407.
- Alexander, G. M., Sherwin, B. B., Bancroft, J., & Davidson, D. W. (1990). Testosterone and sexual behavior in oral contraceptive users and nonusers: a prospective study. *Hormones and Behavior*, *24*, 388–402.
- Archer, J. (2006). Testosterone and human aggression: an evaluation of the challenge hypothesis. *Neuroscience and Biobehavioral Reviews*, *30*, 319–45.
- Bloch, M., Schmidt, P. J., Su, T., Tobin, M. B., & Rubinow, D. R. (1998). Pituitary-adrenal hormones and testosterone across the menstrual cycle in women with premenstrual syndrome and controls. *Biological Psychiatry*, *43*, 897–903.
- Bouissou, M. F. (1990). Effects of estrogen treatment on dominance relationships in cows. *Hormones and Behavior*, *24*, 376–87.
- Burnham, T. C., Chapman, J. F., Gray, P. B., McIntyre, M. H., Lipson, S. F., & Ellison, P. T. (2003). Men in committed, romantic relationships have lower testosterone. *Hormones and Behavior*, *44*, 119–122.
- Cashdan, E. (1995). Hormones. *Sex, and Status in Women*, *29*, 354–366.
- Cobey, K. D., & Buunk, A. P. (2012). Conducting high-quality research on the psychological impact of oral contraceptive use. *Contraception*, *86*, 330–1.
- Cobey, K. D., Pollet, T. V., Roberts, S. C., & Buunk, A. P. (2011). Hormonal birth control use and relationship jealousy: evidence for estrogen dosage effects. *Personality and Individual Differences*, *50*, 315–317.
- Cobey, K. D., Klipping, C., & Buunk, A. P. (2013). Hormonal contraceptive use lowers female intrasexual competition in pair-bonded women. *Evolution and Human Behavior*, *34*, 294–298.
- Dabbs, J. M. (1990). Salivary testosterone measurements: reliability across hours, days, and weeks. *Physiology & Behavior*, *48*, 83–6.
- Darwin, C. (1871). *The descent of man, and selection in relation to sex* (1st ed.). London: John Murray.
- Deady, D. K., Smith, M. J. L., Sharp, M. A., & Al-Dujaili, E. A. S. (2006). Maternal personality and reproductive ambition in women is associated with salivary testosterone levels. *Biological Psychology*, *71*, 29–32.
- Denson, T. F., Mehta, P. H., & Ho Tan, D. (2013). Endogenous testosterone and cortisol jointly influence reactive aggression in women. *Psychoneuroendocrinology*, *38*, 416–24.
- Edelstein, R. S., Chopik, W. J., & Kean, E. L. (2011). Sexuality moderates the association between testosterone and relationship status in men and women. *Hormones and Behavior*, *60*, 248–55.
- Edwards, D. A., & Casto, K. V. (2013). Women's intercollegiate athletic competition: cortisol, testosterone, and the dual-hormone hypothesis as it relates to status among teammates. *Hormones and Behaviour*, *64*, 153–160.
- Gill, J. A. N. (2000). The effects of moderate alcohol consumption on female hormone levels and reproductive function. *Alcohol & Alcoholism*, *35*, 417–423.
- Grant, V. J., & France, J. T. (2001). Dominance and testosterone in women. *Biological Psychology*, *58*, 41–7.

- Gray, P. B., Chapman, J. F., Burnham, T. C., McIntyre, M. H., Lipson, S. F., & Ellison, P. T. (2004). Human male pair bonding and testosterone. *Human Nature, 15*, 119–131.
- Harding, C. F. (1983). Hormonal influences on avian aggressive behavior. In B. Svare (Ed.), *Hormones and aggressive behaviour* (pp. 435–367). New York: Plenum.
- Johnson, S. G., Joplin, G. F., & Burrin, J. M. (1987). Direct assay for testosterone in saliva: relationship with a direct serum free testosterone assay. *Clinica Chimica Acta, 163*, 309–318.
- Kivlighan, K. T., Granger, D. A., & Booth, A. (2005). Gender differences in testosterone and cortisol response to competition. *Psychoneuroendocrinology, 30*, 58–71.
- Kraemer, W. J., & Ratamess, N. A. (2005). Hormonal responses and adaptations to resistance exercise and training. *Sports Medicine, 35*, 339–61.
- Kuzawa, C. W., Gettler, L. T., Huang, Y., & McDade, T. W. (2010). Mothers have lower testosterone than non-mothers: evidence from the Philippines. *Hormones and Behavior, 57*, 441–447.
- Liening, S. H., Stanton, S. J., Saini, E. K., & Schultheiss, O. C. (2010). Salivary testosterone, cortisol, and progesterone: two-week stability, interhormone correlations, and effects of time of day, menstrual cycle, and oral contraceptive use on steroid hormone levels. *Physiology & Behavior, 99*, 8–16.
- Maestripieri, D., Klimczuk, A. C. E., Senenzko, M., Traficante, D. M., & Wilson, M. C. (2013). Relationship status and relationship instability, by not dominance, predict individual differences in baseline cortisol levels. *PLoS ONE, 8*(12), e84003.
- Magee, J. O. E. C., & Galinsky, A. (2008). Social Hierarchy: the self-reinforcing nature of power and status. *Academy of Management Annals, 2*, 1–79.
- McIntyre, M., Gangestad, S. W., Gray, P. B., Chapman, J. F., Burnham, T. C., O'Rourke, M. T., & Thornhill, R. (2006). Romantic involvement often reduces men's testosterone levels—but not always: the moderating role of extrapair sexual interest. *Journal of Personality and Social Psychology, 91*, 642–51.
- Mehta, P. H., & Josephs, R. A. (2010). Testosterone and cortisol jointly regulate dominance: evidence for a dual-hormone hypothesis. *Hormones and Behavior, 58*, 898–906.
- Morris, N. M., Udry, J. R., Khan-Dawood, F., & Dawood, M. Y. (1987). Marital sex frequency and midcycle female testosterone. *Archives of Sexual Behavior, 16*, 27–37.
- Muller, M. N., & Wrangham, R. W. (2004). Dominance, aggression and testosterone in wild chimpanzees: a test of the “challenge hypothesis.” *Animal Behaviour, 67*, 113–123.
- Neave, N., Laing, S., Fink, B., & Manning, J. T. (2003). Second to fourth digit ratio, testosterone and perceived male dominance. *Proceedings of the Royal Society B: Biological Sciences, 270*, 2167–72.
- Piccoli, V., Cobey, K. D., & Carnaghi, A. (2014). Hormonal contraceptive use and the objectification of women and men. *Personality and Individual Differences, 66*, 44–47.
- Pollet, T. V., van der Meij, L., Cobey, K. D., & Buunk, A. P. (2011). Testosterone levels and their associations with lifetime number of opposite sex partners and remarriage in a large sample of American elderly men and women. *Hormones and Behavior, 60*, 72–7.
- Ponzi, D., Klimczuk, A. C. E., Traficante, D. M., & Maestripieri, D. (2015). Perceived dominance in young heterosexual couples in relation to sex, context, and frequency of arguing. *Evolutionary Behavioural Sciences, 1*, 43–54.
- Purifoy, F. E., & Koopmans, L. H. (1979). Androstenedione, testosterone, and free testosterone concentration in women of various occupations. *Biodemography and Social Biology, 26*, 179–188.
- Rada, R. T., Kellner, R., & Winslow, W. W. (1976). Plasma testosterone and aggressive behaviour. *Psychosomatics, 17*, 138–142.
- Roberts, S. C., Cobey, K. D., Klapilová, K., & Havlíček, J. (2014). Oral contraception and romantic relationships: From the lab to the real world. *Human Ethology Bulletin, 29*, 4–13.
- Rose, R. M., Holaday, J. W., & Bernstein, I. S. (1971). Plasma testosterone, dominance rank and aggressive behaviour in male rhesus monkeys. *Nature, 231*, 366–368.
- Schultheiss, O. C., Dargel, A., & Rohde, W. (2003). Implicit motives and gonadal steroid hormones: effects of menstrual cycle phase, oral contraceptive use, and relationship status. *Hormones and Behavior, 43*, 293–301.
- Searcy, W. A., & Wingfield, J. C. (1980). The effects of androgen and antiandrogen on dominance and aggressiveness in male red-winged blackbirds. *Hormones and Behavior, 14*, 126–35.
- Stanton, S. J., & Schultheiss, O. C. (2007). Basal and dynamic relationships between implicit power motivation and estradiol in women. *Hormones and Behavior, 52*, 571–80.
- Swaddle, J. P., & Reiersen, G. W. (2002). Testosterone increases perceived dominance but not attractiveness in human males. *Proceedings of the Royal Society B: Biological Sciences, 269*, 2285–9.
- Udry, J. R., & Talbert, L. M. (1988). Sex hormone effects on personality at puberty. *Journal of Personality and Social Psychology, 54*, 291–295.
- Van Anders, S. M., & Goldey, K. L. (2010). Testosterone and partnering are linked via relationship status for women and “relationship orientation” for men. *Hormones and Behavior, 58*, 820–6.

- Van Anders, S. M., & Watson, N. V. (2006). Relationship status and testosterone in North American heterosexual and non-heterosexual men and women: cross-sectional and longitudinal data. *Psychoneuroendocrinology*, *31*(6), 715–23.
- Van Anders, S. M., & Watson, N. V. (2007). Testosterone levels in women and men who are single, in long-distance relationships, or same-city relationships. *Hormones and Behavior*, *51*(2), 286–91.
- Vittek, J., L’Hommedieu, D. G., Gordon, G. G., Rappaport, S. C., & Louis Southren, A. (1985). Direct radioimmunoassay (RIA) of salivary testosterone: correlation with free and total serum testosterone. *Life Sciences*, *37*(8), 711–716.
- Welling, L. L. M., Jones, B. C., DeBruine, L. M., Conway, C. A., Law Smith, M. J., Little, A. C., & Al-Dujaili, E. A. S. (2007). Raised salivary testosterone in women is associated with increased attraction to masculine faces. *Hormones and Behavior*, *52*(2), 156–61.
- Welling, L. L. M., Puts, D. A., Roberts, S. C., Little, A. C., & Burriss, R. P. (2012). Hormonal contraceptive use and mate retention behavior in women and their male partners. *Hormones and Behavior*, *61*(1), 114–20.
- Wingfield, J. C., Hegner, R. E., Duffy, A. M., Jr., & Ball, G. F. (1990). The “challenge hypothesis”: theoretical implications for patterns of testosterone secretion, mating systems, and breeding strategies. *American Naturalist*, *136*, 829–846.
- Zimmerman, Y., Eijkemans, M., Coelingh Bennink, H. J. T., Blankensin, M., & Fauser, B. (2013a). The effect of combined oral contraception on testosterone levels in healthy women: A systematic review and meta-analysis. *Human Reproduction Update*, *20*, 76–105.
- Zimmerman, Y., Eijkemans, M. J. C., Coelingh Bennink, H. J. T., Blankenstein, M. A., & Fauser, B. C. J. M. (2013b). The effect of combined oral contraception on testosterone levels in healthy women: a systematic review and meta-analysis. *Human Reproduction Update*, *20*(1), 76–105.
- Zimmerman, Y., Foidart, J.-M., Pintiaux, A., Minon, J.-M., Fauser, B. C. J. M., Cobey, K. D., & Coelingh Bennink, H. J. T. (2014). Restoring testosterone levels by adding dehydroepiandrosterone to a drospirenone containing combined oral contraceptive: I endocrine effects. *Contraception*, *91*(2), 127–133.